

RESEARCH ARTICLE

UPPER CRETACEOUS-OLIGOCENE TETHYAN AGGLUTINATED BENTHIC FORAMINIFERAL SPECIES OF THE GENUS *ARENOBULIMINA*

Haidar Salim Anan

Emeritus professor of stratigraphy and paleontology, former Vice president of Al Azhar University-Gaza, P. O. Box 1126, Palestine.

*Corresponding Author Email: profanan@gmail.com

This is an open access journal distributed under the Creative Commons Attribution License CC BY 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

ARTICLE DETAILS

Article History:

Received 03 June 2023
Revised 06 August 2023
Accepted 10 September 2023
Available online 13 September 2023

ABSTRACT

The agglutinated test of the genus *Arenobulimina* has trochospirally enrolled, four or more chambers per whorl, aperture interiomarginal with a simple arch or loop, which has a comprising various species. Seven small Upper Cretaceous-Oligocene benthic foraminiferal species of this agglutinated genus *Arenobulimina* are common in some Tethyan localities: North Atlantic (USA, Trinidad, Labrador Sea), Europe (Spain, France, Germany, Czech, Poland, Sweden), Northeast Africa (Egypt), and Southwest Asia (Palestine, UAE). These species are: Upper Cretaceous *A. americana* Cushman, Late Maastrichtian *A. beitjebrenensis* Anan and *A. jerusalemensis* Anan, Upper Cretaceous-Paleocene *A. truncata* (Reuss), Paleocene *Arenobulimina aegyptiaca* (Said & Kenawy) and Paleocene-Eocene *A. dorbignyi* (Reuss). One Early Eocene - Early-Oligocene species *Arenobulimina kaminskii* is believed here to be new species.

KEYWORDS

Palaeontology, stratigraphy, paleogeography, benthic foraminifera, *Arenobulimina*, Tethys

1. INTRODUCTION

The genus *Arenobulimina* has received much attention from many authors, e.g. (Reuss, 1845 (Germany); Cushman, 1936 (USA); Said & Kenawy, 1956 (Egypt); Gawor-Biedowa, 1969 (Poland), Price, 1977 (NW Europe); Anan, 2022 (Palestine). This study deals with some other Upper Cretaceous-Early Oligocene species of *Arenobulimina* in the Tethys, which have wide geographic distributed in North America, West Europe, North Africa, and

SW Asia (Figure 1). Evolutionary changes of the identified species are indicated by such criteria, such as changes in the test-size, chambers arrangement, type of sutures, periphery or surface ornamentation. These changes help to define the major faunal change through the Maastrichtian-Oligocene times, and can be used in the biostratigraphic subdivisions and paleogeographic and environmental correlations.



Figure 1: Location map of the members of the genus *Arenobulimina* in the Northern Tethys (Spain, France, Germany, Poland, Czech, Sweden) and Southern Tethys (Egypt, Palestine, UAE).

2. MATERIAL AND METHOD

Seven small Upper Cretaceous-Oligocene benthic foraminiferal species of the agglutinated genus *Arenobulimina* are recorded from many localities in the Northern and Southern Tethys. The present study aims at throwing

light on: 1) present together many data scattered in the literature for this genus under a unifying theme, 2) detect its paleontology, stratigraphy and paleogeographic distribution of the different species of that genus, and 3) present a new species *Arenobulimina kaminskii*.

Quick Response Code



Access this article online

Website:
www.geologicalbehavior.com

DOI:
10.26480/gbr.02.2023.70.74

3. TAXONOMY

The members of the genus *Arenobulimina* are characterized by its trochospirally enrolled agglutinated test, four or more chambers per whorl, a simple arch or loop interiomarginal aperture, but without apertural tooth. The taxonomy of Kaminski is followed in this study for diagnostic seven agglutinated foraminiferal benthic specie were erected from six countries in the Tethys (Kaminski, 2014). The new taxonomic considerations are used for many recorded species. These species are illustrated in Plate 1.




Plate 1: Figure 1. *Arenobulimina aegyptiaca* Said & Kenawy (1956) x 30, 2. *A. americana* Cushman (1936) x 45, 3. *A. beitjebrinensis* Anan (2022) x 35, 4. *A. dorbignyi* (Reuss, 1845), x 65, 5. *A. jerusalemensis* Anan (2022), x 35, 6. *A. kaminskii* Anan, n. sp. x 70, 7. *A. truncata* (Reuss, 1845) x 20.

Order Loftusiida Kaminski & Mikhalevich, 2004

Suborder Ataxophragmiina Fursenko, 1958

Genus *Arenobulimina* Cushman, 1927

***Arenobulimina aegyptiaca* Said & Kenawy, 1956**, p. 126, pl. 1, fig. 42 (illustrated specimen) (Plate 1, figure 1)

Remarks: This Paleocene species is characterized by its smooth conical test, tapering at the base, and greatest breadth at the apertural end, aperture a narrow loop-shaped opening at the base of the apertural face. It was recorded from the Paleocene of Sinai (Egypt), and Jabal Malaqet (UAE) (see Figure 1).

***Arenobulimina americana* Cushman, 1936**, p. 27, pl. 4, fig. 9 (Plate 1, figure 2)

Remarks: This Upper Cretaceous species differs from the Paleocene *A. aegyptiaca* by its finely wall arenaceous rather smoothly finished test, and small loop-shaped aperture. It was recorded, so far from USA, Mexico, Brazil and Argentina (Figure 2).



Figure 2: Location map of USA and Mexico (North America) and Brazil and Argentina (South America).

***Arenobulimina beitjebrinensis* Anan, 2022**, p. 27, pl. 1, fig. 58; pl. 2, fig. 3 (= *Arenobulimina* sp. A. Almogi-Labin et al., 1990, p. 578, pl. 2, fig. 5 (Plate 1, figure 3)

Remarks: This Late Maastrichtian species differs from the other members of the genus *Arenobulimina* by its more tapering initial part and more inflated last chambers in the last whorl. It is recorded, so far, from Palestine (Figure 3).



Figure 3: Location map of the two studied sections in Palestine (Jerusalem and Beit Jebrin), as well as Giddi and Nekhl sections, Sinai of Egypt (after Said & Kenawy, 1956).

***Arenobulimina dorbignyi* (Reuss, 1845)** (= *Bulimina dorbignyi* Reuss, 1845, p. 38, pl. 13, fig. 74; Sztrákos, 2005, p. 184, pl. 2, fig. 6) (Plate 1, figure 4)

Remarks: This Paleocene-Eocene species differs from the previous species by its rounded initial part and finely-medium agglutinated grains of the test. It is recorded from Trinidad, Sweden, France, Czech, Poland (see Figure 1).

***Arenobulimina jerusalemensis* Anan, 2022**, p. 26, pl. 1, fig. 59, pl. 2, fig. 4. (= *Arenobulimina* sp. B. Almogi-Labin et al., 1990, p. 578, pl. 2, fig. 4) (Plate 1, figure 5)

Remarks: This Late Maastrichtian species differs from *A. beitjebrinensis* in its shorter length test. It is recorded, so far, from Palestine.

***Arenobulimina kaminskii* Anan, n. sp.** (= *Arenobulimina* sp. Kaminski et al., 1989, p. 730, pl. 6, fig. 4) (Plate 1, figure 6)

Holotype: Illustrated specimen in Plate 1, figure 6.

Diameter: Length 0.25 mm, width 0.20 mm.

Etymology: After Prof. M. Kaminski, King Fahd University of Petroleum and Minerals, Saudi Arabia.

Type locality: Southern Labrador Sea (Figure 4).

Stratigraphic level: Early Eocene-Early Oligocene.

Diagnosis: This long ranged species has trochospirally enrolled test, coarsely agglutinated wall, distinct inflated chambers, sutures slightly depressed, aperture an interiomarginal highly open looped on the apertural face, and without lip or apertural tooth.

Remarks: This new species differs from other recorded species of *Arenobulimina* in its coarsely arenaceous wall, highly and wide opening aperture on the apertural face.




Figure 4: Location map of the studied wells in the Labrador Sea, North Atlantic Ocean.

***Arenobulimina truncata* (Reuss, 1845)** (= *Bulimina (pasternakia) truncata* Reuss, 1845, p. 38, pl. 8, fig. 73) (Plate 1, figure 7)

Remarks: The Upper Cretaceous-Paleocene species distinguished by its tapering initial part with acute angle to the axis of coiling. It has a wide geographic distribution in North America (USA, Mexico, Trinidad) and Europe (Spain, Germany, Czech).

4. STRATIGRAPHIC VALUE OF THE SEVEN BENTHIC FORAMINIFERAL SPECIES OF ARENOBULIMINA

Seven species have wide stratigraphic time from the Upper Cretaceous to Early Oligocene and recorded from many parts of the Tethys: North America to southwest Asia via Europe and north Africa. The stratigraphic value of these taxa are arranged from descending order (older to younger): Upper Cretaceous *Arenobulimina americana* Cushman, Late

Maastrichtian *A. beitjebrinensis* and *A. jerusalemensis* both of Anan, Upper Cretaceous-Paleocene *A. truncata* (Reuss), Paleocene *A. aegyptiaca* Said & Kenawy, Paleocene-Eocene *A. dorbignyi* (Reuss) and Early Eocene-Early Oligocene *A. kaminskii* n. sp. The Upper Cretaceous and Paleocene fauna (*A. americana*, *beitjebrinensis*, *A. jerusalemensis*, *A. truncata* and *A. aegyptiaca*) have mainly a tapering initial part with acute angle to the axis of coiling, and fine to medium-grained agglutinated wall, while the post Paleocene to Oligocene species (*A. dorbignyi* and *A. kaminskii*) have rounded initial part, and coarse-grained agglutinated wall (Figure 5).




Figure 5: The stratigraphic levels of the identified Pre-Maastrichtian-Oligocene species of the genus *Arenobulimina* from the different localities in the Tethys: 1. *Arenobulimina americana* Cushman, 2. *A. beitjebrinensis* Anan, 3. *A. jerusalemensis* Anan, 4. *A. truncata* (Reuss), 5. *A. aegyptiaca* Said & Kenawy, 6. *A. dorbignyi* (Reuss), 7. *A. kaminskii* n. sp

5. PALEOGEOGRAPHY

The Tethys had been connected from west Atlantic Ocean to east with the Pacific Ocean via the Mediterranean Sea during the Cretaceous-Paleogene time. The identified seven species have wide stratigraphic time from the

Upper Cretaceous to Early Oligocene and recorded from many parts of the Tethys: North America (USA, Mexico, Trinidad), South America (Brazil, Argentina), Europe (Spain, Germany, Czech), North Africa (Egypt) and southwest Asia (Palestine, UAE) (Figure 6).



Figure 6: The paleogeographic map at K/T boundary showing the connected seas from west (USA, Atlantic Ocean, Brazil, Argentina) to east (Indian and Pacific Oceans) via the Mediterranean Sea, as well as the locations of the Northern Tethys localities (Spain, France, Germany) and Southern Tethys localities (Egypt, Palestine, UAE) (after Solakius et al., 1990).

According to many authors (e.g. Anan, 1995; Haq and Aubry, 1978; Rosenbaum et al., 2002; Finger, 2013; Anan, 2022). The wide geographic distribution of the identified species proved that the Tethyan realm had been connected with the Atlantic Ocean from west to the Indo-Pacific Ocean to the east, via Mediterranean Sea, crossing the Middle East region during the Paleogene time.

Table (1) shows the paleogeographic distribution of the identified seven agglutinated benthic foraminiferal species in the Upper Cretaceous-Early Oligocene. Some remarks are presented:

1. *Arenobulimina dorbignyi* has wide geographic distribution, which were recorded from five countries: Trinidad, Sweden, France, Czech and Poland.

- 2. *A. truncata* has also wide geographic distribution: USA, Mexico, Spain, Czech.
- 3. *A. americana* also has wide geographic distribution: USA, Mexico, Chile and Argentina.
- 4. *A. aegyptiaca* was recorded from two countries: Egypt and UAE.
- 5. The other three species of *Arenobulimina* was confined to their original type localities: Palestine (*A. beitjebrinensis*, *A. jerusalemensis*), and Labrador Sea (*A. kaminskii*).
- 6. Regardless of unrealized synonymies, this clearly indicates that many species did not have wide geographic or temporal ranges.

Table 1: Paleogeographic distribution of the seven agglutinated benthic foraminifera species of the genus *Arenobulimina* in some Tethyan localities: 1. USA, 2. Mexico, 3. Trinidad, 4. Brazil, 5. Argentina, 6. Labrador Sea, 7 Sweden, 8. Spain, 9. France, 10. Czech, 11. Poland, 12. Egypt, 13. Palestine, 14. UAE (Sp. No. = Species number, x=recorded, - not recorded).

| Sp. No. | countries | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 |
|---------|----------------------|------------------------|---|---|---|---|---|---|---|---|---|----|----|----|----|----|
| | species | | | | | | | | | | | | | | | |
| 1 | <i>Arenobulimina</i> | <i>aegyptiaca</i> | - | - | - | - | - | - | - | - | - | - | - | x | - | x |
| 2 | | <i>americana</i> | x | x | - | x | x | - | - | - | - | - | - | - | - | - |
| 3 | | <i>beitjebrenensis</i> | - | - | - | - | - | - | - | - | - | - | - | - | x | - |
| 4 | | <i>dorbignyi</i> | - | - | x | - | - | - | x | x | x | x | - | - | - | - |
| 5 | | <i>jerusalemensis</i> | - | - | - | - | - | - | - | - | - | - | - | - | x | - |
| 6 | | <i>kaminskii</i> | - | - | - | - | - | x | - | - | - | - | - | - | - | - |
| 7 | | <i>truncata</i> | x | x | - | - | x | - | - | x | - | x | - | - | - | - |

6. PALEOENVIRONMENT

The identified members of the genus *Arenobulimina* includes finely wall arenaceous smoothly finished and also coarsely agglutinated wall tests. According to different authors, the smaller agglutinating foraminifera occur in all marine environments, from marginal to deep, and some are tolerant of hyposalinity as well as normal marine salinity, and/or of hypoxia or dysoxia (e.g. Kaminski et al, 1989; Cetean et al, 2011). Increasingly poorer preservation of agglutinated foraminifers reflects the first appearance of cool, nutrient-poor deep water. During the sea-level maxima calcareous-cemented species such as *Clavulinoides*, *Dorothia*, *Marssonella*, *Arenobulimina*, and *Gaudryina* increase in number, becoming almost equal in numbers with the tubular forms.

7. CONCLUSIONS

The present study deals with the recording of seven identified species of the agglutinated foraminifera genus *Arenobulimina* in fifteen localities of the Tethys: North America (USA, Mexico, Trinidad, Labrador Sea), South America (Brazil, Argentina), Europe (Spain, France, Germany, Sweden, Poland, Czech), Southern Tethys (Egypt, Palestine and UAE). The Tethyan realm had been connected with the Atlantic Ocean from west to the Indo-Pacific Ocean to the east, via Mediterranean Sea, crossing the Middle East region during the Maastrichtian-Paleogene time. Evolutionary changes of the identified species are presented and these changes help to define the major faunal change through the Maastrichtian-Oligocene times, and can be used in the biostratigraphic subdivisions and environmental conditions.

ACKNOWLEDGMENT

The author would like to express his sincere appreciation to the editor of GBR for kind cooperation. I am also indebted to my daughter Dr. Huda Anan for help in the development of the figures, table and plate.

REFERENCES

- Alegret, L., Ortiz, S., and Molina, E., 2009. Extinction and recovery of benthic foraminifera across the Paleocene–Eocene Thermal Maximum at the Alamedilla section (Southern Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 279, Pp. 186-200.
- Almogi-Labin, A., Bein, A., and Sass, E., 1990. Agglutinated foraminifera in organic-rich neritic carbonates (Upper Cretaceous, Israel) and their use in identifying oxygen levels in oxygen-poor environments. *Paleoecology, Biostratigraphy, Paleoceanography and taxonomy of agglutinated Foraminifera*. Kluwer Academic Publication, Netherland, Pp. 565-585.
- Anan, H.S., 1993. Paleocene benthonic foraminifera of Jabal Malaqet, Al Ain region, United Arab Emirates. *Al-Azhar Bulletin of Sci., Al Azhar Univ., Cairo*, 4 (1), Pp. 293-320.
- Anan, H.S., 1995. Late Eocene biostratigraphy of Jabals Malaqet and Mundassa of Al Ain region, United Arab Emirates. *Revue de Micropaléontologie*, 38 (1), Pp. 3-14.
- Anan, H.S., 2022. Contribution to the paleontology, stratigraphy and paleogeography of ninety-seven Southern Tethyan agglutinated foraminiferal species. *Earth and Planetary Science*, 1 (1), Pp. 22-34.

Brotzen, F., 1936. Foraminiferen aus dem schwedischen untersten Senon von Eriksdal in Schonen: *Sveriges Geologiska Undersökning*, C. 396, Arsbok 30 (3).

Cetean, C.G., Bălc, R., Kaminski, M.A., Filipescu S., 2011. Integrated biostratigraphy and Palaeoenvironments of an upper Santonian e upper Campanian succession from the southern part of the Eastern Carpathians, Romania. *Cretaceous Research*, 32, Pp. 575-590.

Cushman, J.A., 1936. New genera and species of the families Verneulinidae and Vulvulinidae and of the subfamily Virguliniinae. *Cushman Foundation for Foraminiferal Research, Special Publication*, 6, Pp.1-71.

Cushman, J.A., 1937. A monograph of the subfamily Virguliniinae of the foraminiferal family Buliminidae. *Special Publications, Cushman Laboratory foraminiferal Research*, 9, Pp.1-228.

Cushman, J.A., Jarvis, P.W., 1932. Upper Cretaceous foraminifera from Trinidad. *United States National Museum Bulletin*, 80 (14), Pp.1-60.

Finger, K., 2013. Miocene foraminifera from the south-central coast of Chile. *Micropaleontology*, 59 (4–5), Pp. 341–492.

Gawor-Biedowa, E., 1969. The genus *Arenobulimina* Cushman from the Upper Albian and Cenomanian of the Polish Lowlands. *Rocznik Polskiego Towarzystwa Geologicznego, Annales De La Societe Geologique De Pologne, Krakow*, 39 (1-3), Pp. 73-104.

Hagn, H., 1953. Die Foraminiferen der Pinswanger Schichten (Unteres Obercampan). Ein Beitrag zur Mikropaläontologie der Helvetischen Oberkreide Südbayerns. *Palaeontographica*, A-104 (1-3), Pp.1-119.

Haq, B.U., Aubry, M.-P., 1978. Early Cenozoic calcareous nannoplankton biostratigraphy and palaeobiogeography of North Africa and the Middle East and Trans-Tethyan correlations. *The Geology of Libya* (1). 2nd Symposium on the geology of Libya, Tripoli, Pp. 271-304.

Kaminski, M.A., Gradstein, F.M., Berggren, W.A., 1989. Paleogene benthic foraminifer biostratigraphy and paleoecology at Site 647, Southern Labrador Sea. *Proceedings of the Ocean Drilling Program, Scientific Results*, 105, Pp.705-730.

Kaminski, M.A., Gradstein, F.M., Berggren, W.A., Geroch, S., Beckmann, J.P., 1988. Flysch-type agglutinated foraminiferal assemblages from Trinidad: Taxonomy, stratigraphy and paleobathymetry. *Geowissenschaftliche Abhandlungen, Bundesanstalt, Vienna*, 41, Pp.155-227.

Neagu, Th., 2014. Turonian-lowermost Emsherian Foraminifera from Dâmbovița Valley area (Upper Basin of the Buzău River-Întorsura Buzăului) - Part II, Benthonic assemblages. *Romanian Journal of Geology*, 58 (1–2), Pp. 27–83.

Price, R.J., 1977. The evolutionary interpretation of the Foraminiferida *Arenobulimina*, *Gavelinella*, and *Hedbergella* in the Albian of North-west Europe. *Palaeontology*, 20 (3), Pp. 503-527.

Reuss, A.E., 1845. Die Versteinerungen der Böhmischen Kreideformation. *Verlagsbuchhandlung E. Schweizerbart'sche, Stuttgart*, 1, Pp.1-58.

Rosenbaum, G., Lister, G.S., Duboz, C., 2002. Relative motions of Africa, Iberia and Europe during Alpine orogeny. *Tectonophysics*, 359, Pp.117-129.

Said, R., Kenawy, A., 1956. Upper Cretaceous and Lower Tertiary foraminifera from northern Sinai, Egypt. *Micropaleontology*, 2 (2),

Pp.105-173.

Solakius, N., Pomoni-Papaioannou, F., Alexopoulos, A., 1990. On the paleogeographic distribution of the Late Maastrichtian planktonic foraminiferal genus *Kassabiana* Salaj & Solakius, 1984. *Acta Geologica Hispanica*, 25 (4), Pp. 289-298.

